

PHANTOM DARK ENERGY AND ITS COSMOLOGICAL CONSEQUENCES

MARIUSZ P. DĄBROWSKI

*Institute of Physics, University of Szczecin,
Wielkopolska 15, 70-451 Szczecin, Poland
mpdabfz@sus.univ.szczecin.pl*

I discuss the dark energy characterized by the violation of the null energy condition ($\varrho + p \geq 0$), dubbed phantom. Amazingly, it is admitted by the current astronomical data from supernovae. We discuss both classical and quantum cosmological models with phantom as a source of matter and present the phenomenon called phantom duality.

1. Introduction

Until a couple of years ago, the standard cosmological source of dark energy was considered to be a slightly negative pressure matter ($-\varrho < p < 0$) with perhaps time-evolving equation of state (quintessence), but not exceeding a “mysterious” barrier $p = -\varrho = -\Lambda$, which corresponded to the cosmological constant. Only the strong energy condition of Hawking and Penrose ($\varrho + 3p > 0$, $\varrho + p > 0$) was presumably violated, and the evolution of the universe in future contained two options: an asymptotic emptiness or a Big-Crunch. However, a deeper analysis of the data from supernovae, cosmic microwave background (WMAP) and large-scale structure (SDSS)¹ shows that the dark energy may also be the matter whose pressure is less than minus the energy density and so violates the null energy condition, and consequently, all the energy conditions. This matter is dubbed phantom,² and it leads to qualitatively new types of the evolution of the universe as a whole. I will discuss these opportunities both in classical³ and quantum⁴ cosmological context.

2. Classical phantom cosmologies - Big-Rip and phantom duality

Phantom is dark energy of a strongly negative pressure which in the easiest case may be simulated by a scalar field ϕ of negative kinetic energy with the Lagrangian

$$L = \frac{l}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) , \quad (1)$$

where $l = -1$ for phantom, $l = +1$ for standard scalar field, and $V(\phi)$ is the potential. In terms of the energy-momentum tensor for a perfect fluid this gives (ϱ - the energy density, p - the pressure): $\varrho = (l/2)\dot{\phi}^2 + V(\phi)$, $p = (l/2)\dot{\phi}^2 - V(\phi)$, and it surely violates the null energy condition $\varrho + p = l\dot{\phi}^2 > 0$, if $l = -1$. There are many other examples where phantom matter appears naturally. For example, this is the case in Brans-Dicke theory in the Einstein frame (provided the Brans-Dicke parameter $\omega < -3/2$), in superstring cosmology, in brane cosmology, and in viscous cosmology. Due to the energy conditions violation, it makes a failure of the standard cosmic censorship conjectures, black hole thermodynamics, positive mass theorems and other renowned theorems of general relativity. It also leads to classical

and quantum instabilities,⁵ which encourages some researchers to disregard it as a serious candidate for the dark energy. However, as it was already mentioned, its main motivation does not come from the theoretical considerations. On the contrary, it comes from the observational data. Of course its stability is a problem, and there are suggestions how to avoid that problem, too.⁶

The most striking result which refers to phantom is that its energy density ϱ grows proportional to the scale factor $a(t)$, i.e.,

$$\varrho \propto a^{3|w+1|} \quad \text{for} \quad w < -1. \quad (2)$$

Then, unlike in a more intuitive standard matter case, where the growth of the energy density corresponds to the decrease of the scale factor, here, the growth of the energy density accompanies the expansion of the Universe. This really gives a new scenario for the future evolution of the universe, which has not been considered so far in cosmology. There is a future singularity, which due to its peculiar properties is called Big-Rip. On the approach to a Big-Rip everything in the universe is pulled apart in a reverse order - first clusters, then galaxies, solar systems, atoms, nuclei etc.²

One of the standard cosmological scenarios in cosmology is the evolution from a Big-Bang to a Big-Crunch. In phantom cosmology it is possible to start with a Big-Rip reach the minimum and terminate at another Big-Rip. This is an example of the phantom duality³ - a new symmetry of the field equations which allows to map a large scale factor (cf. (2)) onto a small one and vice versa due to a change

$$a(t) \leftrightarrow 1/a(t) \quad \text{or} \quad w + 1 \leftrightarrow -(w + 1). \quad (3)$$

Similar symmetry was already discovered in the context of superstring cosmology under the name of the scale factor duality⁷ and further extended onto the brane cosmology⁸ ("phantom triality").

Admission of phantom with $p < -\varrho$ enlarges possible set of cosmological solutions. The most desirable are the solutions which begin with a Big-Bang and terminate at a Big-Rip. This is because they preserve all standard Hot-Big-Bang scenario results and agree with the observational data, which suggests that there was a turning point of the evolution of the universe (the standard matter stopped dominating against phantom) just at the redshift $z = 0.46$.¹ However, as shown explicitly,³ other interesting phantom cosmologies, due to its strongly repulsive contribution to the dynamics appear. Among them there are: Einstein Static Universe with two monotonic solutions - one of them is monotonic towards a Big-Bang and another is monotonic towards a Big-Rip; a monotonic solution which transits between the two Einstein Static Universes (both ways); non-singular oscillating solutions etc.

3. Quantum phantom cosmologies and the large-scale quantum effects

Due to the new types of classical trajectories it is also interesting to ask about possible quantum cosmological implications of phantom dark energy. Despite the

large size of the universe it is advisable to think of quantum effects in the region of Big-Rip singularity since according to (2) the energy density may reach the Planck scale of 10^{19} GeV there. In fact, it was already shown⁹ that there is a dispersion of the wave packets in configuration space (with classical time coordinate eliminated) at Big-Bang, Big-Crunch and at the maximum of expansion point in recollapsing models. The question was studied whether such quantum effects may also appear at a Big-Rip and at the minimum point of expansion.⁴ It emerged that the answer is positive for Big-Rip, but negative so far for the minimum point of expansion. The latter might be due to the simple form of the phantom potentials taken into account, and the question of such quantum effects is still open. Besides, it seems reasonable to ask if spreading of the wave packets in configuration space signaling large-scale quantum effects may appear at the turning point of the expansion - the one for $z = 0.46$ at which standard matter loses domination against the phantom. This is the matter for future work.

4. Discussion

Apart from quintessence ($-\varrho < p < 0$), global acceleration of the universe gives strong observational motivation for a non-standard phantom ($p < -\varrho$) type of matter as a candidate for dark energy. Phantom dark energy may dominate the evolution of the universe and additionally lead it to a Big-Rip singularity - a state in which the whole matter pulled apart, though very dense. A Big-Rip singularity may be dual to a Big-Bang/Big-Crunch singularity. Finally, quantum effects may smear out Big-Rip singularity due to large-scale quantum effects.

Acknowledgments

This research has partially been supported by the Polish Ministry of Science and Education grant No 1P03B 043 29 (years 2005-07).

References

1. J.L. Tonry *et al.*, *Astroph. J.* **594**, 1 (2003); M. Tegmark *et al.*, *Phys. Rev. D* **69**, 103501 (2004).
2. R.R. Caldwell, *Phys. Lett. B* **545**, 23 (2002); R.R. Caldwell, M. Kamionkowski, and N.N. Weinberg, *Phys. Rev. Lett.* **91**, 071301 (2003).
3. M.P. Dąbrowski, T. Stachowiak and M. Szydlowski, *Phys. Rev. D* **68**, 103519 (2003); M.P. Dąbrowski and T. Stachowiak, *Ann. Phys. (N.Y.)* **321**, 771 (2006).
4. M.P. Dąbrowski, C. Kiefer and B. Sandhöfer, *Phys. Rev. D* **74**, 044022 (2006).
5. S.M. Carroll, M. Hofman, and M. Trodden, *Phys. Rev. D* **68**, 023509 (2003); S.D.H. Hsu, A. Jenkins, and M.B. Wise, *Phys. Lett. B* **597**, 270 (2004); J.M. Cline, S. Jeon, and G.D. Moore, *Phys. Rev. D* **70**, 043543 (2004).
6. V.A. Rubakov, hep-th/0604153; P. Creminelli *et al.*, hep-th/0606090.
7. K.A. Meissner and G. Veneziano, *Mod. Phys. Lett. A* **6**, 1721 (1991).
8. J.E. Lidsey, *Phys. Rev. D* **70**, 041302 (2004).
9. C. Kiefer, *Phys. Rev. D* **38**, 1761 (1988); C. Kiefer and H.D. Zeh, *Phys. Rev. D* **51**, 4145 (1995).